



# Measurement of the $\pi\pi$ scattering length from a new structure in the $K \rightarrow 3\pi$ Dalitz plot


**Sergio Giudici**

*Scuola Normale Superiore and INFN Pisa*

**NA48/2 collaboration**

Cambridge, CERN, Chicago, Dubna, Edinburgh, Ferrara, Firenze, Mainz, Northwestern,  
Perugia, Pisa, Saclay, Siegen, Torino, Vienna

# What is the $\pi\pi$ scattering length ?



If  $kR \ll 1$  (small momentum compared to interaction range)

**S-wave** is the dominant component to the total cross section

Bose statistics allows Isospin:  $I = 0, 2$

Scattering matrix

$$S | \pi\pi \rangle = \exp(2i\delta) | \pi\pi \rangle$$

Phases

$$\delta_0 = a_0 k$$

$$\delta_2 = a_2 k$$

-->  $a_0, a_2$  are called Scattering Lengths



# Theoretical predictions

Weinberg (1966)  
Effective field theory for  
strong interaction at low E

$$a_0 m_{\pi^+} = \frac{7m_{\pi^+}^2}{16\pi f_\pi^2} = 0.159$$

$$a_2 m_{\pi^+} = \frac{-m_{\pi^+}^2}{8\pi f_\pi^2} = -0.045$$

Most recently  
Colangelo et al. (2001)  
pt -theory two loops  
*Ref: hep-ph/0103088*

$$a_0 m_{\pi^+} = 0.220 \pm 0.005$$

$$a_2 m_{\pi^+} = -0.0444 \pm 0.0010$$

$$(a_0 - a_2) m_{\pi^+} = 0.265 \pm 0.004$$

- 2% level of accuracy is quite unusual for hadronic physics  
experiments have not yet reached the same level of accuracy



# Experimental Status

1977: measurement by Genève/Saclay experiment @ 20% accuracy

2003: BNL E865 extracts  $a_0$  at 5% accuracy by measuring the form factors of the decay  $K \rightarrow \pi \pi \nu$  with 400,000 events

$$a_0(m) = 0.216 \pm 0.013(\text{stat.}) \pm 0.002(\text{syst.}) \pm 0.002(\text{theor.})$$

*Ref. Pislak et al. (2003) hep-ex/0301040*


Running: Cern experiment DIRAC with a sophisticated technique aims to measure the pionium lifetime @ 10% accuracy

$$\tau \sim 40 \cdot (a_0 - a_2)^2 \cdot 10^{-15} \text{ sec}$$

Pionium is the atom-like electromagnetic bound state ( $\pi^+ \pi^-$ )

# What does it matter to Kaons ?

A diagram contributing to  $K^+ \rightarrow \pi^+ \pi^0 \pi^0$




- Strong rescattering + charge exchange with effective coupling constant  $(a_0 - a_2)$
- Small Pionium formation also expected

# $(a_0 - a_2)$ in $K^+ \rightarrow \pi^+ \pi^0 \pi^0$ decays

Two processes contribute to  $K^+ \rightarrow \pi^+ \pi^0 \pi^0$

- 1) Direct emission of  $\pi^+ \pi^0 \pi^0$
- 2)  $\pi^0 \pi^0$  produced in charged pions rescattering

(1)



$$\mathcal{M}_0 = 1 + gu/2$$


$$u = 2m_K(m_K/3 - E_{\text{odd}}^*)/m_\pi^2$$

Dalitz variable

$$G = 0.638 \pm 0.020$$

(present PDG value)

(2)



$$\mathcal{M}_1 \propto (a_0 - a_2)$$

$$d\Gamma/dm_{\pi\pi} \propto |\mathcal{M}_0 + \mathcal{M}_1|^2$$

Interference is expected

# Cusp-like effect in $K^+ \rightarrow \pi^+ \pi^0 \pi^0$

*Nicola Cabibbo (2004), hep-ph/0405001*

One loop calculation predicts

a cusp located at  $m_{\pi\pi} = 2m_{\pi^+}$

Where the looping  $\pi^+ \pi^-$  pair flips from off-mass to on mass shell


(Remind:  $m_{\pi^0} = 0.135$  GeV ,  $m_{\pi^+} = 0.139$  GeV)

Data collected by NA48 experiment stimulated theoreticians to develop a two loops calculation


*N. Cabibbo and G. Isidori hep-ph/0502130*

$$\partial\Gamma/\partial(m_{\pi\pi})^2 = \mathcal{F}(a_0 - a_2, a_2, g, h, m_{\pi\pi})$$

Indicating a strategy to extract the Scattering lengths from the experimental  $m_{\pi\pi}$  spectrum



# NA48/2 data $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$



The cusp was first seen by I. Mannelli who was looking for the Pionium and N. Cabibbo realized that it was a clean and beautiful example of a general cusp-like behaviour of cross sections next to threshold for new channels (Wigner 1948)




# Why does NA48/2 collect so many $K^\pm$ ?

Direct CP -violation program

1997-2001 NA48/1  $\text{Re}(\epsilon'/\epsilon) = (14.7 \pm 2.2) \times 10^{-4}$  Neutral Kaon system

2003-2004 NA48/2  $\Delta g = (0.5 \pm 3.8) \times 10^{-4}$  Charge Kaon system

Future NA48/3 intent to measure  $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$



$$K^\pm \rightarrow 3\pi$$

$$\Delta\Gamma/du = 1 + gu$$

$$\Delta g = (g^+ - g^-)/(g^+ + g^-)$$


looking for difference in the Dalitz plot distribution

Statistics Collected 2003-2004


$$K^\pm \rightarrow \pi^\pm \pi^0 \pi^0 \sim 2 \cdot 10^8$$

$$K^\pm \rightarrow \pi^\pm \pi^+ \pi^- \sim 4 \cdot 10^9$$

(loose selection cuts)



# NA48 simultaneous unseparated beams



pion decay products stay in pipe

- Split +/-
- Select momentum
- Recombine +/-

- Focusing
- $\mu$  sweeping


- Cleaning
- Beam spectrometer (resolution 0.7 %)

Beams overlap within ~1mm all along the 115m long decay volume (vacuum  $10^{-5}$  mbar)

# $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$ Reconstruction

## Liquid Krypton Calorimeter


photon detection allows to reconstruct the relevant variable  $m_{\pi\pi}$



E resolution  $\sim 1.5\%$ , for  $\langle E \rangle = 10$  GeV  
 position resolution  $\sim 1$  mm  
 time resolution better than 500 ps  
 non linearity  $< 0.1\%$   
 (very stable over 8 years)

$\pi^0 \rightarrow \gamma\gamma$

$\pi^0 \rightarrow \gamma\gamma$




$$\frac{M_{\pi\pi}^2}{4m_{\pi^0}^2} = \frac{\sum_{i,j=1,4} E_i E_j d_{ij}^2}{\left( \sqrt{\sum E_1 E_2 d_{12}^2} + \sqrt{\sum E_3 E_4 d_{34}^2} \right)^2} \quad 1$$

$\pi^\pm \rightarrow \mu^\pm \nu$  makes no effect since charged pion track parameters are not involved in  $m_{\pi\pi}$  computation

2  $\pi^0$  mass constraints  
 Z from 2 vertices average

**Background free**

# Acceptance and $m_{\pi\pi}$ resolution



0.0031  $\text{GeV}^2$  resolution on  $(m_{\pi\pi})^2$  @cusp  
 0.0012  $\div$  0.0120 range elsewhere


$\pi^0\pi^0$  mass constraints and LKR resolution

$$\frac{(E)}{E} = \frac{0.090}{E} \frac{0.032}{\sqrt{E}} \quad 0.0042$$

The acceptance is linearly varying especially around the cusp

The extraction of  $(a_0 - a_2)$  is Montecarlo dependent because of the geometrical acceptance

# Is our Montecarlo good enough ?



Test example:

**Fig. (a)** shows the ratio of the **min  $\gamma$  distance from axis** @ the calorimeter plane between events **above** and **below** the observed cusp.

The Montecarlo well reproduces the data




Fig. (b), (c), (d) show other tests based on other distances

distance (cm)

*a)* min  $\gamma$  distance from axis ; *b)* max  $\gamma$  distance from axis;

*c)* min  $\gamma$ - $\gamma$  distance;      *d)* min  $\gamma$  -  $\pi^\pm$  distance



# Ready to fit the $m_{\pi\pi}$ spectrum

Cabibbo-Isidori 2 loops calculation

$$\partial\Gamma/\partial(m_{\pi\pi})^2 = N \times F(a_0 - a_2, a_2, g, h, m_{\pi\pi})$$

4 free parameters + Norm

Scattering length  $a_0 - a_2$

Scattering length  $a_2$

Linear slope  $g$


Quadratic slope  $h$

The parameter  $g$  and  $h$  are equivalent to the PDG parametrisation of the matrix element for  $\mathbf{K}^\pm \rightarrow \pi^\pm \pi^0 \pi^0$

$$|\mathcal{M}|^2 = (1 + gu + hu^2)$$

Our  $h$  is defined as  $h = h_{\text{PDG}} - g^2/4$

Our Data + MC for acceptance




Data in 420 bins




bin width =  $0.0015 \text{ GeV}^2$  = half the resolution

Cusp located at 50<sup>th</sup> bin

# Fits against various models




The best fit is obtained with the two loops calculation but a small amount of pionium is to be added in bin 50<sup>th</sup> to improve the Chi2



N.B. Cabibbo-Isidori model holds not too far from the cusp --> Not all the spectrum is fitted!

# Fit results



The pionium amount has been fixed according to the prediction

$$\frac{BR(K^+ \rightarrow \pi^+ \text{pionium})}{BR(K^+ \rightarrow 3\pi)} = 0.8 \cdot 10^{-5}$$


*Z.K. Silagadze (1994),  
hep-ph/9411382*

$(a_0 - a_2) \cdot m_\pi$  has low sensitivity to pionium


$$0.1 \cdot \sigma(\text{BR})/\text{BR}$$

Measured by varying the predicted amount in the range  $\pm 50\%$

# Systematics Check: fitting region



Data – Fit



## Choice of the fitting region

systematic on  $(a_0 - a_2) \cdot m_\pi \pm 0.008$

If the excess is interpreted as Pionium formation  
data agree with Silagadze's prediction within  
the (large) error

# Check on photon isolation and Z vertex

Cut on transverse distance @ the calorimeter to avoid photon energy mis-measurement

$$\min(\gamma-\gamma \text{ distance}) > 10 \text{ cm}$$


$$\min(\gamma - \pi^\pm \text{ distance}) > (10 + d) \text{ cm}$$

If  $\pi^\pm$  interacts with Kr, energy may be deposited even at large distance from the impact point

**Default cut  $d=5 \text{ cm}$**  95% of the extra-shower associated to the  $\pi^\pm$  is contained

Trying  $d=5,10,15 \text{ cm}$

$$\text{systematic on } (a_0 - a_2) \cdot m_\pi \pm 0.004$$



Measurement from two different decay regions

$$\text{systematic on } (a_0 - a_2) \cdot m_\pi \pm 0.009$$



## Conclusions on systematics checks

- Excluding ponium from fit region 0.008
- From Cut on track-photon min. distance 0.004
- From dependence on vertex Z coordinate 0.009
- From K<sup>+</sup>/K<sup>-</sup> difference 0.006

**TOTAL (adding in quadrature)**

---

**0.014**



## Preliminary result on $(a_0 - a_2) \cdot m_\pi$

NA48/2 collaboration performed 3 independent analysis on this subject  
two based on a toy (...but not so toy!) Montecarlo  
the third exploiting a professional GEANT-based Montecarlo

The 3 central values found are compatible within an error of 0.001

$$(a_0 - a_2) \cdot m_\pi = 0.281 \pm 0.007 \text{ (stat.)} \pm 0.014 \text{ (syst.)} \pm 0.014 \text{ (theor.)}$$

The theoretical error quoted refers to the Cabibbo-Isidori model and it has been suggested by the authors



# Conclusions


- By including the full 2003 – 04 data sample we expect an increase in statistics from the ~ 30 M events used in the present analysis by a factor of at least 4
  - The study of the systematic uncertainties is well under way, (detailed Monte Carlo, additional systematic effects)
  - The data quality calls for additional theoretical effort in order to extract precise values of the pion – pion scattering parameters
  - A study of the corresponding effects in  $K_L \rightarrow 3\pi^0$  from the 2002 NA48 data has been started
- ... the PDG pages related to the K Dalitz plot description will be deeply modified ...
- Data have also been collected by NA48/2 on  $K+e4$  decays, which will provide independent information on pion – pion scattering



# NA48 Detector


## Main detector components:

- Magnetic spectrometer (4 DCHs):  
redundancy high efficiency;  
 $\Delta p/p = 1.0\% + 0.044\% * p$  [GeV/c]
- Hodoscope  
fast trigger;  
precise time measurement (150ps).
- Liquid Krypton EM calorimeter (LKr)  
High granularity, quasi-homogeneous;  
 $\Delta E/E = 3.2\%/\sqrt{E} + 9\%/E + 0.42\%$  [GeV].
- Hadron calorimeter, muon veto counters, photon vetoes.





# Cusp in $K_L \rightarrow 3\pi^0$



1.5 %

Fitted CUSP position at  $(0.0786 \pm 0.0006) \text{ GeV}^2$

Consistent with the expected value 0.07814