

Searches for Gauge Mediated SUSY Breaking at LEP




Katja Klein

University of Heidelberg / OPAL

XXXVIIth Rencontres de Moriond – EW

March 13th, 2002




- ▷ Introduction
- ▷ Searches for: – topologies with a neutralino NLSP
– topologies with a slepton NLSP
- ▷ Interpretations
- ▷ Summary




All limits are at 95% confidence level.

The LEP Dataset



Distribution of Delivered Integrated luminosity by Energy




Introduction

SUSY particles are not degenerate with Standard Model particles \Rightarrow SUSY is *broken* ...

... either via *gravity* \rightarrow supergravity (SUGRA)

... or via the Standard Model *gauge interactions* \rightarrow **G**auge **M**ediated **S**USY **B**reaking



Advantage of GMSB: gauge interactions are flavour-blind \Rightarrow no FCNC

Six new parameters in the minimal GMSB model:

Λ	SUSY particle mass scale
M	Messenger particle mass scale
N	Messenger index
\sqrt{F}	SUSY breaking scale
$\tan \beta$	Ratio of Higgs vacuum expectation values
$sign(\mu)$	Sign of Higgs sector mixing parameter

GMSB Phenomenology:

SUSY breaking scale $\sqrt{F} \ll 10^{11}$ GeV and $m_{\tilde{G}} \simeq 2.5 \left(\frac{\sqrt{F}}{100 \text{ TeV}} \right)^2 \text{ eV} \rightarrow$ **gravitino \tilde{G} is LSP**
 (LSP = Lightest SUSY Particle)

NLSP (next-to LSP) can be:

- the neutralino: $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$
- the lighter stau: $\tilde{\tau}_1 \rightarrow \tau \tilde{G}$
- all sleptons ("co-NLSP"): $\tilde{\ell}_R \rightarrow \ell \tilde{G}$


The **NLSP decay length** can be *macroscopic*:

$$c\tau \simeq \frac{0.01}{\kappa_\gamma} \left(\frac{100 \text{ GeV}}{m_{\text{NLSP}}} \right)^5 \left(\frac{\sqrt{F}}{100 \text{ TeV}} \right)^4 \text{ cm}$$

(κ_γ = photino component of the $\tilde{\chi}_1^0$; $\kappa_\gamma = 1$ for $\tilde{\ell}$)





\Rightarrow Expect a variety of challenging topologies depending on the NLSP type and its lifetime!

NLSP Type, $\text{sign}(\mu) > 0$




(OPAL GMSB scan)

Topologies in the Neutralino NLSP Scenario:




- ▷ **Prompt neutralino decays:** photons + \cancel{E} (+ leptons / jets)
- ▷ **Intermediate neutralino lifetime:** non-pointing photons + \cancel{E} (+ leptons / jets)
- ▷ **Stable neutralinos:** invisible (or leptons / jets + \cancel{E})

Search for Acoplanar Photons




Zero lifetime \Rightarrow




Search for two high-energetic photons and missing p_T

$130 \leq E_{\text{CM}} \leq 208 \text{ GeV}$
ALEPH DELPHI L3 OPAL

Irreducible background:




Discriminant: recoil mass against the $\gamma\gamma$ system




No excess over BG expectation

→ cross-section limits:



$$M_{\tilde{\chi}_1^0} > 96 \text{ GeV for } M_{\tilde{e}_R} = 2.0 M_{\tilde{\chi}_1^0} \approx M_{\tilde{e}_L}$$




⇒ GMSB interpretation of the famous $e e \gamma \gamma \cancel{E}_T$ event from CDF excluded!
 $(\tilde{e}^+ \tilde{e}^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0 e^+ e^- \rightarrow \gamma \gamma \tilde{G} \tilde{G} e^+ e^-)$

Neutralinos with Intermediate Lifetimes

For **longer** neutralino lifetimes: photons do not point to the primary vertex


- search for *single photons* with large impact parameters ($d_0 > 40$ cm)
- high *calorimeter granularity* allows to reconstruct impact parameter



Main BG: cosmic ray events & detector noise

ALEPH, 189–209 GeV: 2 obs. / 1.0 exp.

DELPHI, 130–209 GeV: 16 obs. / 14.6 exp.




Search for Long-Lived Neutralinos

$\tilde{\chi}_1^0$ stable $\Rightarrow e^+e^- \rightarrow \tilde{\chi}_1^0\tilde{\chi}_1^0$ is invisible \Rightarrow indirect production used to constrain $M(\tilde{\chi}_1^0)$

$\triangleright e^+e^- \rightarrow \tilde{\ell}^+\tilde{\ell}^- \rightarrow \ell^+\tilde{\chi}_1^0\ell^-\tilde{\chi}_1^0$


$\triangleright e^+e^- \rightarrow \tilde{\chi}_1^+\tilde{\chi}_1^- \rightarrow W^{+*}\tilde{\chi}_1^0W^{-*}\tilde{\chi}_1^0$



\rightarrow SUGRA ($\tilde{\chi}_1^0$ is LSP) searches can be used





$\Rightarrow M(\tilde{\chi}_1^0) > 54 \text{ GeV}$ for all lifetimes

($N = 1, \tan \beta = 3, \Lambda = 39 \text{ TeV}, M = 10^{10} \text{ GeV}$)






(theoretical framework: Ambrosanio, Kribs, Martin)

Topologies in the Slepton NLSP Scenario:




- ▷ **Prompt slepton decays:** 2 / 4 / 6 leptons + \cancel{E}
- ▷ **Intermediate slepton lifetime:** tracks with large impact parameters / kinks (+ 2 / 4 leptons)
- ▷ **Stable sleptons:** sleptons (+ 2 / 4 leptons)

Search for Acoplanar Leptons



Search for Sleptons with Intermediate Lifetime

A simulated $\tilde{\mu}^+ \tilde{\mu}^-$ event:



If the slepton decays

- before reaching the tracking chamber:
 - tracks with large impact parameters (d_0)
- in the tracking chamber:
 - tracks with kinks

- BG: – cosmic ray events
 – hadronic interaction with detector material
 – beam related BG

Large d_0 + kinks,
 189–209 GeV:

- A: 1 obs. / 1.1 exp.
 D: 9 obs. / 7.4 exp.
 O: 2 obs. / 2.8 exp. ($\tilde{e}, \tilde{\mu}$)
 7 obs. / 4.4 exp. ($\tilde{\tau}$)

Search for Long-Lived Sleptons

Pair-produced sleptons decay outside detector



- two back-to-back tracks with
- anomalous high or low ionization energy loss (dE/dx)
- no calorimeter signals ("heavy muons")

SM background-free & very high efficiency (OPAL > 90%)




$\sqrt{s} = 189 - 209 \text{ GeV}$:

ALEPH: 1 obs. / 2.3 exp.
 DELPHI: 1 obs. / 2.7 exp.
 OPAL: 1 obs. / 0.6 exp.

OPAL



Combination of
all lifetimes:
Cross-section limits






Combination of all lifetimes: *Slepton mass limits*

$$M_{\tilde{e}} > 66.0 \text{ GeV}$$


$$M_{\tilde{\mu}} > 95.2 \text{ GeV}$$

$$M_{\tilde{\tau}} > 86.1 \text{ GeV}$$


(exp. cross-sections based on Dimopoulos, Thomas, Wells)



$\tilde{\chi}_1^0 \tilde{\chi}_1^0$ Production with a $\tilde{\ell}$ NLSP




→ stau NLSP: $\tilde{\tau}\tilde{\tau}$
 → slepton co-NLSP:
 $\tilde{e}\tilde{e}, \tilde{\mu}\tilde{\mu}, \tilde{\tau}\tilde{\tau}, \tilde{e}\tilde{\mu}, \tilde{e}\tilde{\tau}, \tilde{\mu}\tilde{\tau}$



zero lifetime
topology:

... but also large d_0 & kinks!



E.g. $\tilde{\tau}\tilde{\tau}$, 189–209 GeV:

4 leptons + \cancel{E} :

- A: 22 obs. / 16.5 exp.
- D: 15 obs. / 11.6 exp.
- O: 5 obs. / 5.0 exp.

Large d_0 and kinks:



- A: 5 obs. / 5.3 exp.
- O: 5 obs. / 5.4 exp.

Interpretations in the Minimal GMSB Model

Scan the
GMSB parameters, e.g.:

$$\begin{aligned}
 1 &\leq N \leq 5 \\
 1.5 &\leq \tan \beta \leq 40 \\
 10 \text{ TeV} &\leq M \leq 10^9 \text{ TeV} \\
 \text{sign}(\mu) &= \pm 1 \\
 1 \text{ TeV} &\leq \Lambda \leq \min(\sqrt{F}, M)
 \end{aligned}$$

Use MSSM Higgs exclusion
in $M_h - \sin^2(\beta - \alpha)$ plane



$\Lambda > 10 \text{ (16) TeV}$
(sparticle masses $\propto \Lambda$)

(theoretical framework: Ambrosanio, Kribs, Martin)

Summary

Searches have been performed at LEP for (almost) all topologies predicted within the minimal GMSB model.

No evidence for GMSB has been found at LEP.

Constraints on the GMSB parameter space have been substantially improved by the LEP results.

More results expected: final analyses of all experiments, final LEP combinations, even new channels.

We hope for exciting results from the Tevatron & LHC!