


ep-Physics at High Q^2

Recent Results and Future Perspectives
Testing QCD and Electroweak Theory at HERA

Hans-Christian Schultz-Coulon
Universität Dortmund
[for the H1 and ZEUS collaboration]


Les Arcs, March 2002

HERA Kinematics




NC DIS Event

[as seen in a typical HERA detector]



$$Q^2 = 16950 \text{ GeV}^2$$



Neutral Current DIS Cross Section

$$\frac{d^2\sigma^{NC}(e^\pm)}{dx dQ^2} = \frac{2\pi\alpha^2}{xQ^4} [Y_+ \tilde{F}_2 \mp Y_- x\tilde{F}_3 - y^2 F_L]$$

$$\{Y_\pm = (1 \pm (1-y)^2)\}$$

Helicity structure

F_L : influence small
related to gluon density
[contribution only @ high y]
[LO: $F_L=0$]

LO picture:

$$\tilde{F}_2 = x \sum_i [q_i(x, Q^2) + \bar{q}_i(x, Q^2)] \cdot \mathbf{A}_i$$

$$x\tilde{F}_3 = x \sum_i [q_i(x, Q^2) - \bar{q}_i(x, Q^2)] \cdot \mathbf{B}_i$$


Hard process
[electroweak couplings & propagator]


Proton structure
[fitted in NLO]


$$A_i = Q_q^2 - 2Q_q v_e v_q \chi_Z + Q_q v_e v_q \chi_Z + (v_e^2 + a_e^2)(v_q^2 + a_q^2)(\chi_Z)^2$$

$$B_i = -2Q_q a_e a_q \chi_Z + 4v_e a_e v_q a_q (\chi_Z)^2$$

$$\chi_Z = \frac{1}{4s_W^2 c_W^2} \left(\frac{Q^2}{Q^2 + M_Z^2} \right)$$







NC 'reduced'
Cross Section



$$\tilde{\sigma}_{NC}^{\pm} \equiv \frac{1}{Y_+} \frac{xQ^4}{2\pi\alpha^2} \frac{d^2\sigma^{NC}(e^{\pm})}{dx dQ^2}$$

$$\tilde{\sigma}_{NC}^{\pm} = \tilde{F}_2 \mp f(y)\tilde{F}_3 + g(y)F_L$$

Extraction of:

- $x\tilde{F}_3$ (@ large Q^2)
- quark densities


F_L Extraction Method



- Substantial F_L contribution only at low x i.e. @ high $y = Q^2/xs$
- Subtraction method taking F_2 from QCD Fit (at low y region)

$$F_L \equiv \frac{Y_+}{y^2} [F_2 - A \tilde{\sigma}_{NC}]$$

Normalization factor taken from data with $y < 0.6$ [negligible contribution of F_L]



$x\tilde{F}_3$ Extraction Method

[Using NC e^+p and e^-p cross section]

$$\tilde{\sigma}^{NC}(e^-) = \frac{1}{Y_+} [Y_+ \tilde{F}_2 + Y_- x\tilde{F}_3 - y^2 F_L]$$

$$\tilde{\sigma}^{NC}(e^+) = \frac{1}{Y_+} [Y_+ \tilde{F}_2 - Y_- x\tilde{F}_3 - y^2 F_L]$$


$$x\tilde{F}_3 = \frac{Y_+}{Y_-} \cdot [\tilde{\sigma}^{NC}(e^-) - \tilde{\sigma}^{NC}(e^+)]$$

sensitivity to
valence quark densities

$$x\tilde{F}_3 \sim [q(x, Q^2) - \bar{q}(x, Q^2)]$$

sensitive only @ high Q^2
where γZ interference is sizeable

additional factor needed if
 e^+p and e^-p data taken at different beam energies



$x\tilde{F}_3$

- rises with Q^2 (@ fixed x) due to propagator


$$\chi_Z = \frac{1}{4s_W^2 c_W^2} \left(\frac{Q^2}{Q^2 + M_Z^2} \right)$$

- agreement of data with prediction from QCD fit

contribution
to $x\tilde{F}_3$: <3%

$$x\tilde{F}_3 = Q_e a_e \underbrace{\{2Q_q a_q x [q_i - \bar{q}_i]\}}_{\sim xF_3^{\gamma Z}} \cdot \chi_Z + 2v_e a_e \{2v_q a_q x [q_i - \bar{q}_i]\} \cdot (\chi_Z)^2$$

$$\sim Q_e a_e \{xF_3^{\gamma Z}\} \cdot \chi_Z$$



$x F_3^{\gamma Z}$

- Q^2 dependence (from scaling violation) expected to be small
- direct comparison of measurements at different Q^2 possible


$$\int_0^1 F_3^{\gamma Z} = \int_0^1 2Q_q a_q [q_i - \bar{q}_i] = 2Q_u a_u N_u + 2Q_d a_d N_d = \frac{5}{3} \cdot (1 - \alpha_s/\pi)$$

[sum rule a la Gross Llewellyn-Smith]


H1 measurement: $\int_{0.02}^{0.65} F_3^{\gamma Z} = 1.88 \pm 0.44$

H1 QCD Fit: $\int_{0.02}^{0.65} F_3^{\gamma Z} = 1.11$

} agreement within 2 standard deviations



Charged Current Cross Section




$$\frac{d^2\sigma^{CC}(e^-)}{dx dQ^2} = \frac{\pi\alpha^2}{4s_W^2} \frac{1}{(Q^2 + M_W^2)^2} [\underbrace{u + c}_{\text{Probes u-quark density}} - (1-y)^2(\bar{d} + \bar{s})]$$

$$\frac{d^2\sigma^{CC}(e^+)}{dx dQ^2} = \frac{\pi\alpha^2}{4s_W^2} \frac{1}{(Q^2 + M_W^2)^2} [\bar{u} + \bar{c} - (1-y)^2(\underbrace{d + s}_{\text{Probes d-quark density}})]$$

CC DIS Event

[as seen by the other typical HERA detector]





Reduced CC Cross Section

[Sensitivity to u/d quark densities]

$$\tilde{\sigma}_{CC}^{\pm} = \left[\frac{4s_W^2 (Q^2 + M_W^2)^2}{\pi\alpha^2} \right] \frac{d\sigma^{CC}(e^{\pm})}{dx dQ^2}$$


$$\tilde{\sigma}_{CC}^{-} = x[u + c + (1-y)^2(\bar{d} + \bar{s})]$$

$$\tilde{\sigma}_{CC}^{+} = x[\bar{u} + \bar{c} + (1-y)^2(d + s)]$$



Knowledge of d/u Ratio

d/u



Fit to

- HERA ep data [H1 1994, ZEUS 1994]
- Fixed Target proton and deuteron data [E665, NMC, BCDMS, SLAC]
- Neutrino data [CCFR]
- Drell-Yan data [E866]


Large uncertainty at high x
[dependence on parameterisation]
[e.g. due to nuclear corrections]

Can be further constrained
with NC and CC HERA data.

Valence Quark Distribution

@ high x

H1 Preliminary



NLO QCD fit

using high Q^2 ,
neutral/charged current,
 e^+p and e^-p data.


Quark densities

determined via local
extraction method for
data points where the
 xq_v contribution is $>70\%$.


$$xq_v = \sigma_{meas} \cdot \left(\frac{xq_v}{\sigma} \right)_{fit}$$

More statistics needed
to constrain behaviour of d_v , u_v further.

ZEUS Microvertex Detector





HERA Upgrade



- ZEUS:**
- Lumi System
 - Triggering
 - Microvertex
 - Forw. Tracking

- H1:**
- Lumi System
 - Triggering
 - Fwd/Bwd Silicon
 - Forw. Tracking

PostUpgrade Luminosity




	Peak spec. Lumi [$\text{cm}^{-2}\text{s}^{-1}\text{mA}^{-2}$]	Peak Lumi [$\text{cm}^{-2}\text{s}^{-1}$]	int. Lumi [pb^{-1}/y]
2000	0.7×10^{30} [$I_e=50 \text{ mA}, I_p=100 \text{ mA}, \#\text{Bunches} \sim 200$]	1.7×10^{31}	70
PostUpgr.	1.5×10^{30} [$I_e=60 \text{ mA}, I_p=140 \text{ mA}, \#\text{Bunches} \sim 200$]	6×10^{31}	240

New beam optics with stronger focusing provides expected improvement


Polarization for H1 and ZEUS

- utilises Compton scattering
- measures energy weighted asymmetry




- utilises Compton scattering
- measures spatial asymmetry

Utilising Polarisation



$$\tilde{\sigma}_{CC}^{\pm} = (1 \pm P)\tilde{\sigma}_{CC, (P=0)}^{\pm}$$

+: Probe d_v quark distribution ($P=+1$)
 -: Probe u_v quark distribution ($P=-1$)




$$\begin{aligned} \tilde{\sigma}_{NC}^{\pm} &= \tilde{\sigma}_{NC,0}^{\pm} + P\tilde{\sigma}_{NC,P}^{\pm} \\ &= f(q, \bar{q}, EW \text{ couplings}) \end{aligned}$$

Four independent equations
 one each for $Q_e = \pm 1$ and $P = \pm 1$.

Possibility to

Disentangle individual quark densities
 Measure EW couplings v_u, v_d, a_u, a_d

LHC parton kinematics




HERA @ high x , high Q^2
 + Fixed Target

Test of QCD evolution
 over 4 order of magnitude.

important for LHC and
 e.g. the prediction of
 Higgs/W cross sections.

EW Couplings v_u, a_u



Conclusions

Neutral and charged current cross section
consistent with SM

Electroweak effects used as tool to extract
proton structure @ high x

HERA II:

Test of QCD evolution up to highest Q^2
Constrain valence quark distributions at high x
Determine EW couplings v_u, a_u, v_d, a_d