
$e^+e^- \rightarrow 3$ jets and event shapes at NNLO

Aude Gehrmann-De Ridder

in collaboration with T. Gehrmann, E.W.N. Glover, G. Heinrich

XLIII th Rencontres de Moriond, "QCD and High Energy Interactions"

La Thuile, 12.03.08

$e^+e^- \rightarrow 3 \text{ jets and event shapes}$

Classical QCD observable

- testing ground for QCD: **perturbation theory, power corrections and logarithmic resummation**
- precision measurement of strong coupling constant α_s
- current error on α_s from jet observables dominated by theoretical uncertainty:
S. Bethke, 2006

$$\alpha_s(M_Z) = 0.121 \pm 0.001(\text{experiment}) \pm 0.005(\text{theory})$$

- theoretical uncertainty largely from missing higher orders
- current status: **NLO plus NLL resummation**

Jets in Perturbation Theory

Jet Description

- Partons are combined into jets using the same jet algorithm as in experiment

LO	NLO	NNLO
		
each parton forms 1 jet on its own	2 partons in 1 jet, 1 parton experimentally unresolved	3 partons in 1 jet, 2 partons experimentally unresolved

Current state-of-the-art: NLO plus resummation of all-order logarithms (NLLA)


Need for higher orders:

- reduce error on α_s
- better matching of parton level and hadron level

Ingredients to NNLO $e^+e^- \rightarrow 3\text{-jet}$

Two-loop matrix elements

$|\mathcal{M}|^2_{2\text{-loop}, 3 \text{ partons}}$




explicit infrared poles from loop integrals

L. Garland, T. Gehrmann, N. Glover, A. Koukoutsakis,
E. Remiddi; S. Moch, P. Uwer, S. Weinzierl

One-loop matrix elements

$|\mathcal{M}|^2_{1\text{-loop}, 4 \text{ partons}}$




explicit infrared poles from loop integral and
implicit infrared poles due to single unresolved radiation

Z. Bern, L. Dixon, D. Kosower, S. Weinzierl;
J. Campbell, D.J. Miller, E.W.N. Glover

Tree level matrix elements

$|\mathcal{M}|^2_{\text{tree}, 5 \text{ partons}}$



implicit infrared poles due to double unresolved radiation

K. Hagiwara, D. Zeppenfeld;
F.A. Berends, W.T. Giele, H. Kuijf;
N. Falck, D. Graudenz, G. Kramer

Infrared Poles cancel in the sum

NLO Subtraction

Structure of NLO m -jet cross section (subtraction formalism):

Z. Kunszt, D. Soper

$$d\sigma_{NLO} = \int_{d\Phi_{m+1}} \left(d\sigma_{NLO}^R - d\sigma_{NLO}^S \right) + \left[\int_{d\Phi_{m+1}} d\sigma_{NLO}^S + \int_{d\Phi_m} d\sigma_{NLO}^V \right]$$

- $d\sigma_{NLO}^S$: local counter term for $d\sigma_{NLO}^R$
- $d\sigma_{NLO}^R - d\sigma_{NLO}^S$: free of divergences, can be integrated numerically

General methods at NLO

- Dipole subtraction
S. Catani, M. Seymour; NNLO: S. Weinzierl
- \mathcal{E} -prescription
S. Frixione, Z. Kunszt, A. Signer;
NNLO: S. Frixione, M. Grazzini; V. Del Duca, G. Somogyi, Z. Trocsanyi
- Antenna subtraction (derived from physical matrix elements)
D. Kosower; J. Campbell, M. Cullen, N. Glover;
NNLO: T Gehrmann, E.W.N. Glover, AG

Numerical Implementation

Parton-level event generator

Starting point: $e^+e^- \rightarrow 4$ jets at NLO (EERAD2: J. Campbell, M. Cullen, N. Glover)

- contains already 4-parton and 5-parton matrix elements
- is based on NLO antenna subtraction

additions:

- NNLO subtraction terms (5-parton channel)
- 1-loop-single unresolved integrated subtraction term (4-parton channel)
- 2-loop matrix element (3-parton channel)

checks:


- analytic cancellation of infrared poles
- local cancellations along phase space trajectories approaching singular limits
- confirmation of our result by independent calculation of all logarithmically enhanced terms in the thrust distribution using the SCET formalism
very recently: T. Becher, M. Schwartz

Three-jet cross section at NNLO

NNLO corrections: jet rates

Three-jet fraction in Durham jet algorithm

$$y_{i,j,D} = \frac{2 \min(E_i^2, E_j^2) (1 - \cos \theta_{ij})}{E_{vis}^2}$$



- NNLO corrections small
- substantial reduction of scale dependence
- better description towards lower jet resolution
- need: resummation at low y_{cut} and hadronization corrections

Event shapes variables

Standard Set of LEP

- Thrust (E. Farhi)

$$T = \max_{\vec{n}} \left(\frac{\sum_{i=1}^n |\vec{p}_i \cdot \vec{n}|}{\sum_{i=1}^n |\vec{p}_i|} \right)$$

- Heavy jet mass (L. Clavelli, D. Wyler)

$$\rho = M_i^2 / s = \frac{1}{E_{\text{vis}}^2} \left(\sum_{k \in H_i} |\vec{p}_k| \right)^2$$

- C-parameter: eigenvalues of the tensor (G. Parisi)

$$\Theta^{\alpha\beta} = \frac{1}{\sum_k |\vec{p}_k|} \frac{\sum_k p_k^\alpha p_k^\beta}{\sum_k |\vec{p}_k|}$$


- Jet broadenings (S. Catani, G. Turnock, B. Webber)

$$B_i = \left(\frac{\sum_{k \in H_i} |\vec{p}_k \times \vec{n}_T|}{2 \sum_k |\vec{p}_k|} \right)$$

$$B_W = \max(B_1, B_2) \quad B_T = B_1 + B_2$$


- $3j \rightarrow 2j$ transition parameter in Durham algorithm y_{23}^D

S.Catani, Y.L.Dokshitzer, M.Olsson, G.Turnock, B.Webber



Event shapes at NNLO


NNLO thrust and heavy mass distributions



- NNLO corrections sizeable, non uniform: 15-20% in T , 10% in ρ
- theory uncertainty reduced by about 40 %
- large $1 - T, \rho > 0.33$: kinematically forbidden at LO
- small $1 - T, \rho$: two-jet region, need matching onto NLL resummation
Just completed: T. Gehrmann, G. Luisoni, H. Stenzel
- need to include hadronization corrections

Event shapes at NNLO+NLLA


NNLO+NLLA thrust and heavy mass



- (NNLO +NLLA) compared to (NNLO) prediction
 - slightly better description towards the 2-jet limit
 - In the 3-jet region, two predictions in agreement
 - further improvement needed: by including hadronization corrections


Comparison with data

High precision data from all LEP experiments, compare here to ALEPH



- include quark mass effects to NLO
P. Nason, C. Oleari
W. Bernreuther, A. Brandenburg, P. Uwer
G. Rodrigo, A. Santamaria
- include hadronization corrections
HERWIG: B. Webber et al.
ARIADNE: T. Sjostrand et al.
- try new fit of α_s , based on ALEPH analysis
G. Dissertori, T. Gehrmann,
G. Heinrich, H. Stenzel, AG

Extraction of α_s



- scale uncertainty reduced by factor 2 compared to NLO; factor 1.3 compared to NLLA+NLO
- scatter among values from different observables reduced very substantially at NNLO
→ genuine NNLO effect

Result for all ALEPH event shapes of LEP1/LEP2

$$\alpha_s(M_Z) = 0.1240 \pm 0.0008(stat) \pm 0.0010(exp) \pm 0.0011(had) \pm 0.0029(theo)$$

Summary and Conclusions

- Completed the calculation of the NNLO corrections to $e^+e^- \rightarrow 3$ jets using antenna subtraction method
- Presented results for the 3-jet cross section (Durham algorithm)
 - improvement towards lower y_{cut}
 - reduced scale dependence
- Presented results for event shapes in e^+e^- annihilation
 - size of the NNLO corrections sizeable but not uniform
 - improved theoretical uncertainty
 - considerably better consistency between observables
 - new NNLO extraction of α_s
- new result for NLLA+NNLO corrections for event shapes
 - better description of 2-jet limit compared to NNLO
 - NNLO and NNLO+NLLA effects in agreement in the 3-jet region
- next steps
 - α_s from NLLA+NNLO
 - moments of event shapes and analytical power corrections


Back-up slides

Colour structure of NNLO 3-jet

Decomposition into leading and subleading colour terms

$$\sigma_{NNLO} = (N^2 - 1) \left[N^2 A_{NNLO} + B_{NNLO} + \frac{1}{N^2} C_{NNLO} + NN_F D_{NNLO} \right. \\ \left. + \frac{N_F}{N} E_{NNLO} + N_F^2 F_{NNLO} + N_{F,\gamma} \left(\frac{4}{N} - N \right) G_{NNLO} \right]$$

- last term: closed quark loop coupling to vector boson, numerically tiny



$$N_{F,\gamma} = \frac{\left(\sum_q e_q \right)^2}{\sum_q e_q^2}$$

- most subleading colour: C_{NNLO} , E_{NNLO} , F_{NNLO} , (G_{NNLO})
QED-type contributions: gluons \rightarrow photons
- simplest term: F_{NNLO} , only 3 parton and 4 parton contributions

Colour-ordered antenna functions


Antenna Functions

- colour-ordered pair of hard partons (radiators) with radiation in between
 - hard quark-antiquark pair
 - hard quark-gluon pair
 - hard gluon-gluon pair
- three-parton antenna \longrightarrow one unresolved parton
- four-parton antenna \longrightarrow two unresolved partons
- can be at tree level or at one loop
- all three-parton and four-parton antenna functions can be derived from physical matrix elements, normalised to two-parton matrix elements
 - $q\bar{q}$ from $\gamma^* \rightarrow q\bar{q} + X$
 - qg from $\tilde{\chi} \rightarrow \tilde{g}g + X$
 - gg from $H \rightarrow gg + X$

Antenna functions

Quark-antiquark

consider subleading colour (gluons photon-like)




$$|M_{q\bar{q}ggg}|^2(1, 3, 4, 5, 2) \xrightarrow{1||3} |M_{q\bar{q}gg}|^2(\widetilde{13}, 4, 5, \widetilde{23}) \times X_{132}$$

with

$$X_{132} = \frac{|M_{q\bar{q}g}|^2}{|M_{q\bar{q}}|^2} \equiv A_3^0(1_q, 3_g, 2_{\bar{q}})$$

Antenna functions


Quark-gluon



$$|M_{q\bar{q}q\bar{q}g}|^2(1, 3, 4, 5, 2) \xrightarrow{3||4} |M_{q\bar{q}gg}|^2(\widetilde{13}, \widetilde{34}, 5, 2) \times X_{134}$$

with hard radiators:

quark ($\widetilde{13}$) and gluon ($\widetilde{34}$)



q^* : spin 1/2, colour triplet

$q(\widetilde{13})$: spin 1/2, colour triplet


$g(\widetilde{34})$: spin 1, colour octet

Off-shell matrix element: violates $SU(3)$ gauge invariance

Antenna functions

Quark-gluon

Construct colour-ordered qg antenna function from $SU(3)$ gauge-invariant decay:
 neutralino \rightarrow gluino + gluon (T. Gehrmann, E.W.N. Glover, AG)




$\tilde{\chi}$: spin 1/2, colour singlet

\tilde{g} : spin 1/2, colour octet

g : spin 1, colour octet

Gluino \tilde{g} mimics quark and antiquark (same Dirac structure), but is octet in colour space



$\tilde{\chi} \rightarrow \tilde{g}g$ described by effective Lagrangian

H. Haber, D. Wyler


$$\mathcal{L}_{\text{int}} = i\eta \bar{\psi}_{\tilde{g}}^a \sigma^{\mu\nu} \psi_{\tilde{\chi}} F_{\mu\nu}^a + (\text{h.c.})$$

Antenna function

$$X_{134} = \frac{|M_{\tilde{g}q'\bar{q}'}|^2}{|M_{\tilde{g}g}|^2} \equiv E_3^0(1_q, 3_{q'}, 4_{\bar{q}'})$$

Antenna functions


Gluon-gluon



$$|M_{q\bar{q}gggg}|^2(1, 3, 4, 5, 2) \xrightarrow{3||4} |M_{q\bar{q}gg}|^2(1, \widetilde{34}, \widetilde{45}, 2) \times X_{345}$$

$H \rightarrow gg$ described by effective Lagrangian

F. Wilczek; M. Shifman, A. Vainshtein, V. Zakharov



$$\mathcal{L}_{\text{int}} = \frac{\lambda}{4} H F_{\mu\nu}^a F_a^{\mu\nu}$$

Antenna function



$$X_{345} = \frac{|M_{ggg}|^2}{|M_{gg}|^2} \equiv F_3^0(3_g, 4_g, 5_g)$$

Antenna functions

	tree level	one loop
<u>quark-antiquark</u>		
$qg\bar{q}$	$A_3^0(q, g, \bar{q})$	$A_3^1(q, g, \bar{q}), \tilde{A}_3^1(q, g, \bar{q}), \hat{A}_3^1(q, g, \bar{q})$
$qgg\bar{q}$	$A_4^0(q, g, g, \bar{q}), \tilde{A}_4^0(q, g, g, \bar{q})$	
$qq'\bar{q}'\bar{q}$	$B_4^0(q, q', \bar{q}', \bar{q})$	
$qqq\bar{q}$	$C_4^0(q, q, \bar{q}, \bar{q})$	
<u>quark-gluon</u>		
qgg	$D_3^0(q, g, g)$	$D_3^1(q, g, g), \hat{D}_3^1(q, g, g)$
$qggg$	$D_4^0(q, g, g, g)$	
$qq'\bar{q}'$	$E_3^0(q, q', \bar{q}')$	$E_3^1(q, q', \bar{q}'), \tilde{E}_3^1(q, q', \bar{q}'), \hat{E}_3^1(q, q', \bar{q}')$
$qq'\bar{q}'g$	$E_4^0(q, q', \bar{q}', g), \tilde{E}_4^0(q, q', \bar{q}', g)$	
<u>gluon-gluon</u>		
ggg	$F_3^0(g, g, g)$	$F_3^1(g, g, g), \hat{F}_3^1(g, g, g)$
$gggg$	$F_4^0(g, g, g, g)$	
$gq\bar{q}$	$G_3^0(g, q, \bar{q})$	$G_3^1(g, q, \bar{q}), \tilde{G}_3^1(g, q, \bar{q}), \hat{G}_3^1(g, q, \bar{q})$
$gq\bar{q}g$	$G_4^0(g, q, \bar{q}, g), \tilde{G}_4^0(g, q, \bar{q}, g)$	
$q\bar{q}q'\bar{q}'$	$H_4^0(q, \bar{q}, q', \bar{q}')$	

Extraction of α_s


Uncertainty from renormalisation scale




Event shapes at NNLO

NNLO corrections: broadenings

wide jet boadening B_W




total jet boadening B_T



- NNLO corrections for B_W smaller than for B_T
- again require matching onto NLL resummation and hadronization corrections
- observe: small corrections for Y_3 ; large corrections for C
- reduction of dependence on renormalisation scale by 30–60%

Three-jet cross section at NNLO

NNLO corrections: jet rates



● substantial improvement towards lower y_{cut}

● two-jet rate now NNNLO