

THE SM AND SUSY AFTER THE 2011 LHC RESULTS

G. ALTARELLI

*Dipartimento di Fisica 'E. Amaldi', Università di Roma Tre and INFN, Sezione di Roma Tre,
I-00146 Rome, Italy*

and

*CERN, Department of Physics, Theory Unit,
CH-1211 Geneva 23, Switzerland*

We present a short review of the LHC results at 7 TeV and their implications on the Standard Model (SM) and on its Supersymmetric (SUSY) extension. In particular we discuss the exclusion range for the SM Higgs mass, the tantalizing hint of an excess at $m_H \sim 125$ GeV, the negative results of searches for SUSY particles (as well as for any other new physics) and the present outlook

1 The first LHC results

The main LHC results so far (with more than $5.5 fb^{-1}$ of integrated luminosity collected by each large experiment at 7 TeV) are listed here, as presented at Moriond 2012.

1) A robust exclusion interval for the SM Higgs has been established which greatly extends the previous range; precisely at 95 % c.l. the excluded intervals of mass are ATLAS: 110 -117.5, 118.5 -122.5, 129 - 539 GeV and CMS: 127.5 - 600 GeV (note that ATLAS also excludes at 95% c.l. a large part of the mass range 110-122.5 GeV, while CMS has some excess in that region). In addition, there is some tantalizing indication for $m_H \sim 125$ GeV. In this respect, what is encouraging is that an excess is seen in the $\gamma\gamma$ mass distribution both in ATLAS (2.8σ at 126.5 GeV) and CMS (2.9σ at 125 GeV). Also there is a hint for $ZZ \rightarrow 4l^\pm$ in ATLAS (2.1σ at 125 GeV: 3 events) and the Tevatron reports a small excess spread over a large interval in $b\bar{b}$ and WW (2.7σ in 115 -135 GeV). These accumulations are all compatible with $m_H \sim 125$ GeV. Further encouragement has been missed because in CMS a possible hint in $ZZ \rightarrow 4l^\pm$ is at a different mass (2.1σ at 119.5 GeV: 3 events) and in ATLAS the number of WW events is less than expected. Overall the evidence for $m_H \sim 125$ GeV could still evaporate and we need to wait for the outcome of the 2012 run, that, with an expected additional integrated luminosity of $15 fb^{-1}$ per experiment at 8 TeV, should either definitely confirm or exclude $m_H \sim 125$ GeV.

2) No evidence of new physics has been found although a big chunk of new territory has been explored.

3) Important results on B and D decays have been obtained mainly by LHCb, whose performance has been exceedingly good (but also on some issues by ATLAS and CMS), e.g. $B_s \rightarrow J\psi\phi$, $B_s \rightarrow \mu\mu$, CP violation in D decay.... Most of the results go in the direction of the SM. For CP violation in D decay, it could indeed be a sign of new physics but, in view of the uncertainty in the SM prediction, it is difficult to be sure.

2 The Higgs Problem

The experimental verification of the Standard Model (SM)[?] cannot be considered complete until the predicted physics of the Higgs sector[?] is not established by experiment. Indeed the Higgs problem is really central in particle physics today[?]. In fact, the Higgs sector is directly related to most of the major open problems of particle physics, like the flavour problem[?] and the hierarchy problem[?], the latter strongly suggesting the need for new physics near the weak scale, which could also clarify the Dark Matter identity.

It is clear that the fact that some sort of Higgs mechanism is at work has already been established. The W and Z longitudinal degrees of freedom are borrowed from the Higgs sector and are an evidence for it. In fact the couplings of quarks and leptons to the weak gauge bosons W^\pm and Z are indeed experimentally found to be precisely those prescribed by the gauge symmetry^{?,?}. To a lesser accuracy the triple gauge vertices γWW and ZWW have also been found in agreement with the specific predictions of the $SU(2) \otimes U(1)$ gauge theory. This means that it has been verified that the gauge symmetry is unbroken in the vertices of the theory: all currents and charges are indeed symmetric. Yet there is obvious evidence that the symmetry is instead badly broken in the masses. The W or the Z with longitudinal polarization that are observed are not present in an unbroken gauge theory (massless spin-1 particles, like the photon, are transversely polarized). Not only the W and the Z have large masses, but the large splitting of, for example, the top-bottom quark doublet shows that even the global weak SU(2) is not at all respected by the fermion spectrum. Symmetric couplings and totally non symmetric spectrum is a clear signal of spontaneous symmetry breaking and its implementation in a gauge theory is via the Higgs mechanism. The big remaining questions are about the nature and the properties of the Higgs particle(s). The LHC has been designed to solve the Higgs problem.

And indeed the SM Higgs is close to be observed or excluded! Either the SM Higgs is very light ($\lesssim 128$ GeV) or rather heavy (i.e. $\gtrsim 600$ GeV). The range $m_H = 122 - 128$ GeV, where possibly there is a signal, is in agreement with precision tests, compatible with the SM (the data are in fair agreement with the SM Higgs cross-sections[?]) and also with the SUSY extensions of the SM. Actually, $m_H \sim 125$ GeV is what one expects from a direct interpretation of EW precision tests: no fancy conspiracy with new physics to fake a light Higgs while the real one is heavy. On the contrary, $m_H \gtrsim 600$ GeV would point to the conspiracy alternative (but no conspirators have been found nearby!). Thus there is really a great suspense on the LHC run this year.

What if the evidence $m_H \sim 125$ GeV evaporates in 2012? Can we do without the Higgs? Suppose we take the gauge symmetric part of the SM and put masses by hand. What is the fatal problem at the LHC scale? The most immediate disease that needs a solution is that in the absence of a Higgs particle or of an alternative mechanism, violations of unitarity appear in scattering amplitudes involving longitudinal gauge bosons (those most directly related to the Higgs sector) at energies in the few TeV range[?]. A crucial question for the LHC is to identify the mechanism that avoids the unitarity violation: is it one or more Higgs bosons or some new vector boson (like additional gauge bosons W' , Z' or Kaluza-Klein recurrences or resonances from a strong sector)? Thus something must happen at the few TeV scale! It is not possible that neither the Higgs nor new physics are present at the Electro-Weak (EW) scale (the only caution is whether the LHC can completely explore the EW scale).

It is well known that there are theoretical bounds on the Higgs mass valid if one assumes that the SM, with only one Higgs doublet, is valid up to a large energy scale Λ where eventually new physics appears. An upper limit on m_H (with mild dependence on m_t and α_s) is obtained, as described in[?], from the requirement that no Landau pole appears, up to the scale Λ , in the Higgs quartic coupling λ , or in simpler terms, that the perturbative description of the theory remains valid up to Λ . The Higgs mass enters because it fixes the initial value of the quartic

Higgs coupling λ in the running from the EW scale up to Λ . Even if Λ is as small as a few TeV the limit is well within the LHC range $m_H < 600 - 800$ GeV and becomes $m_H < 180$ GeV for $\Lambda \sim M_{Pl}$. This upper limit on the Higgs mass in the SM has played a crucial role in the LHC design whose mission requires that the whole allowed range is within reach of the machine. A lower limit on m_H is derived from the requirement of vacuum stability[?], i.e. that the quartic Higgs coupling λ does not turn negative in its running up to Λ (if so the energy would become negative and unbound at large absolute values of the field). Actually, in milder form, one can tolerate a moderate instability, compatible with the present age of the Universe[?]. A recent thorough reanalysis of this issue[?] has concluded that, given the experimental values of m_t and α_s , for $\Lambda \sim M_{GUT} - M_{Pl}$ the stability bound is very close to $m_H = 130$ GeV. The value $m_H \sim 125$ GeV would imply that, in the absence of new physics, our Universe becomes metastable at a scale $\Lambda \sim 10^{10}$ GeV. But the lifetime of our vacuum, for scales up to the Planck mass, would be larger than the age of the Universe. The SM remains viable with some early Universe implications. The vacuum could be stabilized by very little additional new physics (like, for example a heavy singlet S with a large VEV below the metastability scale[?]). Large Majorana neutrino masses can also have an impact on the running[?]. On the basis of this discussion we can conclude that a 125 GeV Higgs would be nearly perfect for a pure and simple SM up to M_{Pl} , just a little bit below the optimal range $130 \lesssim m_H \lesssim 180$ GeV. Incidentally, the possibility that $m_H \sim 130$ GeV, so that the SM becomes unstable precisely at around the Planck mass, and its implications have been studied in the literature[?].

3 Outlook on Avenues beyond the Standard Model

No signal of new physics has been found neither in EW precision tests nor in flavour physics. Given the success of the SM why are we not satisfied with that theory? Why not just find the Higgs particle, for completeness, and declare that particle physics is closed? As well known, the reason is that there are both conceptual problems and phenomenological indications for physics beyond the SM. On the conceptual side the most obvious problems are the proliferation of parameters, the puzzles of family replication and of flavour hierarchies, the fact that quantum gravity is not included in the SM and the related hierarchy problem. Among the main phenomenological hints for new physics we can list the constraints from coupling constant merging in Grand Unified Theories (GUT's), Dark Matter, neutrino masses (explained in terms of L non conservation), baryogenesis and the cosmological vacuum energy (a gigantic naturalness problem). The computable evolution with energy of the effective gauge couplings clearly points (better in SUSY than in the SM) towards the unification of the electro-weak and strong forces at scales of energy $M_{GUT} \sim 10^{15} - 10^{16}$ GeV which are close to the scale of quantum gravity, $M_{Pl} \sim 10^{19}$ GeV. One is led to imagine a unified theory of all interactions also including gravity (at present superstrings provide the best attempt at such a theory). Thus GUT's and the realm of quantum gravity set a very distant energy horizon that modern particle theory cannot ignore. Can the SM without new physics be valid up to such large energies? Indeed, some of the SM problems could be postponed to the more fundamental theory at the Planck mass. For example, the explanation of the three generations of fermions and the understanding of fermion masses and mixing angles can be postponed. But other problems must find their solution in the low energy theory. In particular, the structure of the SM could not naturally explain the relative smallness of the weak scale of mass, set by the Higgs mechanism at $1/\sqrt{G_F} \sim 250$ GeV with G_F being the Fermi coupling constant. This so-called hierarchy problem[?] is due to the instability of the SM with respect to quantum corrections. This is related to the presence of fundamental scalar fields in the theory with quadratic mass divergences and no protective extra symmetry at $\mu = 0$, with μ the scalar mass. For fermion masses, first, the divergences are logarithmic and, second, at $m = 0$ an additional symmetry, i.e. chiral symmetry, is restored. Here, when

talking of divergences, we are not worried of actual infinities. The theory is renormalizable and finite once the dependence on the cut-off Λ is absorbed in a redefinition of masses and couplings. Rather the hierarchy problem is one of naturalness. We can look at the cut-off as a parameterization of our ignorance on the new physics that will modify the theory at large energy scales. Then it is relevant to look at the dependence of physical quantities on the cut-off and to demand that no unexplained enormously accurate cancellations arise.

The hierarchy problem can be put in less abstract terms: loop corrections to the Higgs mass squared are quadratic in the cut-off Λ . The most pressing problem is from the top loop (the heaviest particle, hence the most coupled to the Higgs). If we demand that the correction does not exceed the light Higgs mass indicated by the precision tests, Λ must be close, $\Lambda \sim o(1 \text{ TeV})$. So a crucial question for the LHC to answer is: what damps the top loop contribution? Similar constraints arise from the quadratic Λ dependence of loops with gauge bosons and scalars, which, however, lead to less pressing bounds. So the hierarchy problem demands new physics to be very close. Actually, this new physics must be rather special, because it must be very close, yet its effects are not clearly visible in EW precision tests (the "LEP Paradox"?) now also accompanied by a similar "flavour paradox"?. Examples? of proposed classes of solutions for the hierarchy problem are SUSY, technicolor, "Little Higgs" models, extra dimensions, effective theories for compositeness etc or the alternative, extreme, point of view given by the anthropic solution. In the following, after a comment on the anthropic route, I will discuss the quest for SUSY in some detail, while the alternative solutions to the hierarchy problem will be considered in the companion presentation by Mariano Quiros?.

4 An extreme solution: the anthropic way

The observed value of the cosmological constant Λ poses a tremendous, unsolved naturalness problem?. Yet the value of Λ is close to the Weinberg upper bound for galaxy formation?. Possibly our Universe is just one of infinitely many bubbles (Multiverse) continuously created from the vacuum by quantum fluctuations. Different physics takes place in different Universes according to the multitude of string theory solutions? ($\sim 10^{500}$). Perhaps we live in a very unlikely Universe but the only one that allows our existence??. I find applying the anthropic principle to the SM hierarchy problem somewhat excessive. After all one can find plenty of models that easily reduce the fine tuning from 10^{14} to 10^2 : why make our Universe so terribly unlikely? If to the SM we add, say, supersymmetry, does the Universe become less fit for our existence? In the Multiverse there should be plenty of less fine tuned Universes where more natural solutions are realized and yet are suitable for our living. By comparison the case of the cosmological constant is a lot different: the context is not as fully specified as the for the SM (quantum gravity, string cosmology, branes in extra dimensions, wormholes through different Universes....). While I remain skeptical I would like here to sketch one possibility on how the SM can be extended in agreement with the anthropic idea. If we ignore completely the hierarchy problem and only want to reproduce the most compelling data that demand new physics beyond the SM, a possible scenario is the following one. The SM is to be completed by a light Higgs and no other new physics is in the LHC range (how sad!) except perhaps a Z' , for example a Z'_{B-L} . In particular there is no SUSY in this model. At the GUT scale of $M_{GUT} \gtrsim 10^{16}$ GeV the unifying group is $SO(10)$, broken at an intermediate scale, typically $M_{int} \sim 10^{10} - 10^{12}$ down to a subgroup like the Pati-Salam group $SU(4) \otimes SU(2)_L \otimes SU(2)_R$ or some other one?. Note that unification in $SU(5)$ would not work because we need a group of rank larger than 4 in order to allow for a two step (at least) breaking needed, in the absence of SUSY, to restore coupling unification and to avoid a too fast proton decay. The Dark Matter problem should be solved by axions?. Lepton number violation, Majorana neutrinos and the see-saw mechanism give rise to neutrino mass and mixing. Baryogenesis occurs through leptogenesis?. One should

one day observe proton decay and neutrino-less beta decay. None of the alleged indications for new physics at colliders should survive (in particular even the claimed muon $(g-2)$ discrepancy should be attributed, if not to an experimental problem, to an underestimate of the theoretical errors or, otherwise, to some specific addition to the above model[?]). This model is in line with the non observation of $\mu \rightarrow e\gamma$ at MEG[?], of the electric dipole moment of the neutron[?] etc. It is a very important challenge to experiment to falsify this scenario by establishing a firm evidence of new physics at the LHC or at another "low energy" experiment.

5 Supersymmetry

In the limit of exact boson-fermion symmetry[?] the quadratic divergences of bosons cancel so that only log divergences remain. However, exact SUSY is clearly unrealistic. For approximate SUSY (with soft breaking terms), which is the basis for all practical models, Λ is essentially replaced by the splitting of SUSY multiplets. In particular, the top loop is quenched by partial cancellation with s-top exchange, so, to limit the fine-tuning the s-top cannot be too heavy. The existing limits on SUSY particles (even before the LHC), EW precision tests, success of the Cabibbo-Kobayashi-Maskawa theory of quark mixing and of CP violation, absence of Flavour Changing Neutral Currents, all together, impose sizable fine tuning particularly on minimal realizations. Yet SUSY is a completely specified, consistent, computable model, perturbative up to M_{Pl} . Important phenomenological indications in favour of SUSY are that coupling unification takes place with greater accuracy in SUSY than in the SM and that proton decay bounds are not in contradiction with the predictions. Grand Unification (GUT's) and SUSY go very well together: this is unique among new physics models. Other non standard models^{?,?} (little Higgs, composite Higgs, Higgsless....) all become strongly interacting and non perturbative at a multi-TeV scale. Two Higgs doublets are expected in SUSY[?]. The EW symmetry breaking can be triggered by the H_u mass becoming negative at low energy in the running down from the GUT scale, due to the large top Yukawa coupling. SUSY offers a good Dark Matter candidate: the neutralino (actually more than one candidate, e.g. also the gravitino). In summary SUSY remains the reference model for new physics. But the negative result of the search for SUSY at the LHC, where a big chunk of new territory has been explored in the last year run, has imposed new strong constraints on SUSY models. And the hint of $m_H = 125$ GeV, if confirmed, does even more restrict the parameter space of these models ($m_H = 125$ GeV is a bit too heavy: near the upper bound on m_H in the MSSM).

Even the Minimal SUSY Model (MSSM)[?] has more than 100 parameters (mostly from the SUSY soft breaking terms). Simplified versions with a drastic reduction of parameters are used for practical reasons, e.g. the CMSSM, where C stands for Constrained, or mSUGRA, i.e. minimal SuperGravity (often the two names are confused): with universal gaugino and scalar soft terms at the GUT scale, the set of parameters is drastically reduced down to $m_{1/2}$, m_0 , A_0 (the s-top mixing parameter), $\tan\beta$ and $\text{sign}(\mu)$. Similarly in the Non Universal Higgs Mass models NUHM1,2: masses for H_u , H_d (1 or 2 masses) different from m_0 are added. It is only these oversimplified models that are now cornered. A more flexible setup but, apparently still manageable, is the MSSM with CP and R conservation (pMSSM: p for phenomenological)[?] in terms of 19 parameters (M_A , $\tan\beta$, 3 gaugino masses, 3 mixing parameters A_u , A_d , A_e , μ and 10 s-fermion masses, with degenerate first 2 generations) recently studied in several works.

Many different new physics signatures have been searched at the LHC at 7 TeV with no positive outcome in a variety of channels involving combinations of charged leptons, jets and missing energy. All kinds of models for new physics can be compared with these data, not only SUSY. For SUSY the resulting limits depend on the assumptions on the spectrum, but, in the CMSSM, generically imply that gluinos and degenerate s-quarks are heavier than 500 - 1000 GeV. In addition to these limits the impact of $m_H \sim 125$ GeV on SUSY models is important

? For example, minimal models with gauge mediation or anomaly mediation are disfavoured ? (predict m_H too light) although some versions, like gauge mediation with extra vector like matter ?, could still work. Specific models that give up naturalness but remain predictive like split SUSY or heavy SUSY have seen their allowed domain restricted ?. Gravity mediation ? is in better shape but CMSSM, mSUGRA, NUHM1,2 are only marginally consistent and need s-quarks heavy, A_t large and lead to tension with the muon (g-2). In fact the muon magnetic moment would point to light SUSY, more precisely to light EW gauginos and s-leptons. This type of light SUSY would also improve the EW precision fit (by predicting a heavier m_W than the SM for the experimental value of m_t and a light Higgs). Several groups (for example, see ?) have repeated the fit to EW precision tests in the CMSSM, also including the additional data on the muon ($g-2$), the Dark Matter relic density and rare $b \rightarrow s\gamma$ decay modes. Before the LHC results the promising outcome of this exercise was that the central value of the lightest Higgs mass m_H went up (in better harmony with the bound from direct searches) with moderately large $\tan\beta$ and relatively light SUSY spectrum ?. After the LHC bounds one finds that the best fit Higgs mass is 125 GeV only if the result on the muon (g-2) is removed from the fit, while, with the (g-2) included, the best fit Higgs mass value is 119 GeV. In other words, in the CMSSM there is a sizable tension between the muon (g-2) and $m_H \sim 125$ GeV. Also normally too much Dark Matter is predicted in the CMSSM or mSUGRA for $m_H = 125$ GeV. In comparison, the upper limit on m_H is larger in the pMSSM: $m_H \lesssim 135$ GeV ?.

The problem with SUSY is that one expected its discovery already at LEP2 on the basis of complete naturalness applied to minimal models. With the recent LHC data ever increasing fine tuning appears to be needed in the minimal versions. However less fine tuning is necessary if non minimal models are assumed. One must go beyond the CMSSM, mSUGRA, NUHM1,2. And indeed there is still plenty of room for more sophisticated versions of SUSY as a solution to the hierarchy problem. The simplest new ingredients that are studied at present are either heavy first 2 generations ? and/or an extra Higgs singlet ?.

The first option is still within the MSSM framework. Note that, on the one hand, it is mostly gluinos and 1-2 generation s-quarks that are affected by the LHC limits while EW s-particles and s-tops are less constrained. On the other hand, what is really needed for naturalness in the MSSM ? is that the s-tops (they directly enter at one loop in the radiative corrections to the Higgs mass), their isospin partners the s-bottoms, as well as the lightest higgsino (related to the μ parameter), and also gluinos (that contribute, with a strong coupling, in the radiative corrections at two loops) must be relatively light (below, say, 1 TeV). As remarked already long ago ? an inverted s-quark spectrum with heavier 1st-2nd and lighter 3rd generation s-quarks has several advantages in flavour and CP violation problems. This option has been widely reanalysed recently in the present context. If gluinos are forced to only decay into final states involving s-tops or s-bottoms, their mass limits are considerably less stringent. Similarly the present lower limit on the lightest s-top mass is a few hundred GeV.

By adding an extra singlet Higgs ? one goes beyond the MSSM. In a promising class of models a singlet Higgs S is added with coupling $\lambda SH_u H_d$. The μ term arises from the S Vacuum Expectation Value (VEV) and the μ problem is solved in that the S VEV can naturally be of order of the soft terms that break SUSY. Mixing with S can modify the Higgs mass and couplings at tree level. In particular, the restrictions on the Higgs mass, valid at tree level in the MSSM that demand substantial corrections from loops, can be relaxed (no need of large s-top mixing, less fine tuning). The new coupling λ grows with the scale. If we impose that the theory remains perturbative up to M_{GUT} then we must have $\lambda \lesssim 0.7$. This is the case of the NMSSM (Next to Minimal SSM). For $m_H \sim 125$ GeV larger values of λ allow for lighter s-tops, no large s-top mixing and much less fine tuning. For $\lambda \sim 1-2$ we are in the so-called λ -SUSY regime (for $\lambda \gtrsim 2$ the theory becomes non perturbative already at 10 TeV). The fine tuning can be really reduced to a few percent even with a s-top of mass above 1 TeV. The presence of an extra

Higgs singlet adds one more neutral scalar particle to the spectrum. After symmetry breaking the mixing between S and the doublet Higgs leads to two eigenstates of mass that replace the single lightest Higgs h (for not too large λ a 2 by 2 matrix mixing approximation is valid while for $\lambda \gtrsim 0.7$ the full mixing matrix must be considered). The state at 125 GeV could be the lightest, but it is not excluded that at 125 GeV the heaviest of the two is seen while the lightest escaped detection at LEP[?]. In fact the mixing also modifies the couplings and may be that the lightest eigenstate has suppressed couplings to gauge bosons. In this case the heavier one at 125 GeV would get enhanced couplings to gauge bosons. Indeed there is a tenuous indication that the 125 GeV state may have a slightly enhanced coupling to $\gamma\gamma$.

6 Conclusion

The most exciting result of the 2011 LHC run is that the SM Higgs is close to be observed or excluded! The present, very solid, exclusion ranges for the SM Higgs have much restricted the mass interval for the SM Higgs: either the SM Higgs is very light (115 - 128 GeV) or very heavy (i.e. $\gtrsim 600$ GeV). The range $m_H = 122 - 128$ GeV where some excess is observed is in agreement with precision tests, compatible with the SM and also with the SUSY extensions of the SM. This hint is very exciting but could still disappear with more statistics. Thus the outcome of the 2012 LHC run at 8 TeV is of extreme interest for particle physics.

The search for new physics is the other big issue. No signals have shown up so far in spite of the many channels explored and of the large slice of parameter space that has been for the first time explored. Optimistic expectations of an early success have been deceived. But the LHC experiments are just at the start and larger masses can be reached in 2012 and even more in the 14 TeV phase. Still supersymmetry remains the standard way beyond the SM. It is true that we could have expected the first signals of SUSY already at LEP, based on naturalness arguments applied to the most minimal models like the CMSSM or mSUGRA. But the general MSSM is still very much viable, for example in the versions with heavy 1 - 2 generation s-quarks[?]. Among non minimal models the most studied possibility are based on the addition of an extra singlet S to the Higgs sector[?] (NMSSM and λ - SUSY). The absence of SUSY signals has also stimulated the development of new ideas like those of extra dimensions and composite Higgs (discussed in the talk by M. Quiros[?]). The extreme anthropic proposal that naturalness could be irrelevant for the very particular physics that is valid in our exceptional Universe, just one among many in the Multiverse, is boosted now by the absence of new physics signals at the LHC. Only experiment can choose among these and other possibilities. Supersymmetry? Compositeness? Extra dimensions? Anthropic? We shall see!

I am very grateful to J. Tran Thanh Van and the Organisers of the 2012 Rencontres de Moriond, in particular B. Klima and B. Pietrzyk, for inviting me to give this talk. My related research has been funded by the COFIN program (PRIN 2008), the INFN- Roma Tre, and by the European Commission, under the networks "LHCPHENONET" and "Invisibles".

References

1. A review of the SM can be found in G. Altarelli, in Elementary Particles (Landolt-Brnstein, New Series / Elementary Particles, Nuclei and Atoms), ed. by H. Schopper, Springer, Berlin, 2008, chap. 2, 3 and 4.
2. See, for example, A. Djouadi, Phys. Rept. **457** 1 (2008), ArXiv:hep-ph/0503172.
3. For a pedagogical review, see J.D. Wells, ArXiv:0909.4541.
4. For a review, see G. Isidori, Y. Nir and G. Perez, ArXiv:1002.0900.

5. E. Gildener, Phys. Rev. D 14, 1667 (1976); L. Maiani, in *Vector bosons and Higgs bosons in the Weinberg-Salam theory of weak and electromagnetic interactions. Proc. of the Summer School on Particle Physics*, Gif-sur-Yvette, 3-7 Sep 1979, Ed. by M. Davier *et al.*, IN₂P₃, Paris, France (1979); M. Veltman, Acta Phys. Polon. **B12** (1981) 437; E. Witten, Nucl. Phys. **B 188** (1981) 513 and Phys. Lett. **B 105** (1981) 267.
6. The LEP Electroweak Group, <http://lepewwg.web.cern.ch/LEPEWWG/>.
7. For a review see, for example, G. Altarelli and M. Grunewald, Phys.Rept., 189403 (2004), ArXiv: hep-ph/0404165.
8. J.R. Espinosa *et al*, ArXiv:1202.3697, P.P. Giardino *et al*, ArXiv:1203.4254.
9. B. W. Lee, C. Quigg and H.B. Thacker, Phys.Rev. **D16** 1519 (1977).
10. T. Hambye and K. Riesselmann, Phys.Rev.**D55** 7255 (1997) and references therein.
11. N. Cabibbo, L. Maiani, G. Parisi and R. Petronzio, *Nucl. Phys.* **B 158** (1979) 295; P.Q. Hung, *Phys. Rev. Lett.* **42** (1979) 873; M. Lindner, *Z. Phys.* **C 31** (1986) 295; M. Lindner, M. Sher and H. Zaglauer, *Phys. Lett.* **B 228** (1989) 139. M. Sher, *Phys. Rep.* **179** (1989) 273; B. Schrempp and M. Wimmer, *Prog. Part. Nucl. Phys.* **37** (1996) 1. M. Sher, *Phys. Lett.* **B 317** (1993) 159; *addendum ibid.* **331** (1994) 448; G. Altarelli and G. Isidori, *Phys. Lett.* **B 337** (1994) 141; J.A. Casas, J.R. Espinosa and M. Quirós, *Phys. Lett.* **B 342** (1995) 171.
12. G. Isidori, G. Ridolfi and A. Strumia, Nucl. Phys. **B609387** (2001) and references therein.
13. J. Elias-Mir *et al*, ArXiv:1112.3022.
14. J. Elias-Mir *et al*, ArXiv:1203.0237.
15. C. Wetterich, ArXiv:1112.2910; F. Bezrukov *et al*, ArXiv:1205.2893 and references therein.
16. R. Barbieri and A. Strumia, ArXiv:hep-ph/0007265.
17. For an overview see C. Grojean, ArXiv:0910.4976.
18. M. Quiros, these Proceedings.
19. J. Frieman, M. Turner and D. Huterer, Ann.Rev.Astron.Astrophys. **46** 385 (2008), ArXiv:0803.0982.
20. S. Weinberg, Phys. Rev. Lett. **59**, 2607 (1987).
21. M. R. Douglas, ArXiv:hep-th/0602266; ArXiv:1204.6626.
22. N. Arkani-Hamed and S. Dimopoulos, *JHEP* **0506**, 073 (2005), ArXiv:hep-th/0405159; N. Arkani-Hamed *et al*, Nucl.Phys. **B709** 3 (2005), ArXiv:hep-ph/0409232; G. Giudice and A. Romanino, *Nucl.Phys.* **B699**, 65 (2004), Erratum-*ibid.* **B706**, 65 (2005), ArXiv:hep-ph/0406088; N. Arkani-Hamed, S. Dimopoulos, S. Kachru, ArXiv:hep-ph/0501082; G. Giudice, R. Rattazzi, *Nucl.Phys.* **B757**, 19 (2006), ArXiv:hep-ph/0606105.
23. G. Giudice, in "Perspectives on LHC Physics", ed. by G. Kane and A. Pierce, World Sci., ArXiv:0801.2562.
24. S. Bertolini, L. Di Luzio and M. Malinsky, ArXiv:1205.5637 and references therein.
25. R. D. Peccei and H. R. Quinn, Phys. Rev. D 16 (1977) 1791; R. D. Peccei and H. R. Quinn, Phys. Rev. Lett. 38 (1977) 1440; S. Weinberg, Phys. Rev. Lett. 40 (1978) 223; F. Wilczek, Phys. Rev. Lett. 40 (1978) 279.
26. For a general review, see, for example, W. Buchmuller, R.D. Peccei and T. Yanagida, Ann.Rev.Nucl.Part.Sci.**55**311 (2005), ArXiv:hep-ph/0502169.
27. Muon g-2 Collab., G.W. Bennett *et al.*, Phys. Rev.**D73**072003 (2006).
28. K. Kannike *et al*, ArXiv:1111.2551.
29. The MEG Collaboration, J. Adam *et al*, ArXiv:1107.5547.
30. C. A. Baker *et al*, Phys. Rev. Lett.**97**:131801,2006; ArXiv:hep-ex/0602020v3.
31. For a review, see, for example, A. Hoecker, ArXiv:1201.5093.
32. For recent pedagogical reviews see, for example, S.P. Martin, hep-ph/9709356; I.J. R. Aitchison, ArXiv:hep-ph/0505105; M. Drees, R. Godbole and P. Roy, *Theory and Phenomenology of Sparticles*, World Sci. (2005).

33. A. Djouadi, Phys. Rept. **459** 1 (2008), ArXiv:hep-ph/0503173.
34. A. Djouadi and S. Rosiers-Lees et al., hep-ph/9901246; A. Djouadi, Phys. Rept. 459, 1 (2008), A. Arbey, M. Battaglia and F. Mahmoudi, Eur.Phys.J. C72 (2012) 1847, ArXiv:1110.3726; Eur.Phys.J. C72 (2012) 1906, ArXiv:1112.3032; ArXiv:1205.2557.
35. F. Mahmoudi, ArXiv:1205.3100.
36. A. Arbey et al, Phys.Lett. B708 (2012) 162, ArXiv:1112.3028; P. Draper et al, ArXiv:1112.3068.
37. M. Endo et al, ArXiv:1112.56.53, J. L. Evans et al, ArXiv:1201.2611.
38. G. Giudice and A. Strumia, ArXiv:1108.7067.
39. M. Holthausen, K. S. Lim and M. Lindner, ArXiv:1112.2415; H. Baer, V. Barger and A. Mustafayev, ArXiv:1112.3017; S. Heinemeyer, O. Stal and G. Weiglein, ArXiv:1112.3026; M. Carena et al, ArXiv:1112.3336; O. Buchmueller et al, ArXiv:1112.3564; S. Akula et al, ArXiv:1112.3645; ArXiv:1202.4038; M. Kadastik et al, ArXiv:1112.3647; C. Strege et al, ArXiv:1112.4192; C. Beskidt et al, ArXiv:1202.3366; J. L. Feng and D. Sanford, ArXiv:1205.2372.
40. O. Buchmueller *et al.*, JHEP **0809** 117 (2008), ArXiv:0808.4128; Eur.Phys.J. **C64** 391 (2009), ArXiv:0907.5568.
41. S. Dimopoulos and G. Giudice, Phys.Lett. B357, 573 (1995), ArXiv:hep-ph/9507282; A.Pomarol and D. Tommasini, Nucl.Phys. B466 (1996) 3, ArXiv:hep-ph/9507462; A. G. Cohen, D. Kaplan, and A. Nelson, Phys.Lett. B388, 588 (1996), ArXiv:hep-ph/9607394; R. Barbieri, G.R. Dvali, L. J. Hall; Phys.Lett. B377 (1996) 76, ArXiv:hep-ph/9512388.
42. R. Essig et al, ArXiv:1110.6443; Y. Kats et al, ArXiv:1110.6444; ArXiv:1110.6443; M.Papucci, J.T. Ruderman and A. Weiler, ArXiv:1110.6926; C. Brust et al, ArXiv:1110.6670; C. Csaki, L. Randall and J. Terning, ArXiv:1201.1293; G. Larsen, Y. Nomura and H. L.L. Roberts, ArXiv:1202.6339.
43. R. Barbieri et al, Phys. Rev. D 75, 035007 (2007), ArXiv:hep-ph/0607332; L. Cavicchia, R. Franceschini and V. S. Rychkov, Phys. Rev. D77, 055006 (2008), ArXiv:0710.5750; R. Barbieri et al, JHEP **0803**, 005 (2008), ArXiv:0712.2903; J. Cao and J. M. Yang, Phys. Rev. D78, 115001 (2008), ArXiv:0810.0989; P. Lodone, JHEP **1005**, 068 (2010), ArXiv:1004.1271; R. Barbieri et al, JHEP **1008**, 024 (2010), ArXiv:1004.2256; R. Franceschini and S. Gori, JHEP **1105**, 084 (2011), ArXiv:1005.1070; Y. Nomura, D. Poland and B. Tweedie, Phys. Lett. B 633, 573 (2006), ArXiv:hep-ph/0509244; M. Bastero-Gil et al, Phys. Lett. B 489, 359 (2000), ArXiv:/0006198; A. Delgado, C. Kolda, J. P. Olson and A. de la Puente, Phys. Rev. Lett. 105, 091802 (2010), ArXiv:1005.1282; G. G. Ross and K. Schmidt-Hoberg, ArXiv:1108.1284; U. Ellwanger and C. Hugonie, Mod. Phys. Lett. A 22, 1581 (2007), ArXiv:hep-ph/0612133; U. Ellwanger, C. Hugonie and A. M. Teixeira, Phys. Rept. 496, 1-77 (2010), ArXiv:0910.1785; L. J. Hall, D. Pinner and J. T. Ruderman, ArXiv:1112.2703; S. F. King, M. Muhlleitner and R. Nevzorov, ArXiv:1201.2671.
44. U. Ellwanger, ArXiv:1112.3548.